

The ScanMars radar onboard AMADEE-18 analog mission to Mars

Alessandro Frigeri (1), Maurizio Ercoli (2), Cristina Pauselli (2), Gernot Groemer (3)

(1) Istituto di Astrofisica e Planetologia Spaziali, INAF, Rome, Italy (Alessandro.Frigeri@iaps.inaf.it, Phone:+39-06-4993-4227); (2) Università degli Studi di Perugia, Perugia, Italy; (3) Österreichisches Weltraum Forum, Innsbruck, Austria.

Abstract

Simulated, or analogue, planetary missions are putting the base for the forthcoming extraterrestrial explorations, testing new technologies and refining existing operative procedures [4, 2, 3].

In February 2018, the Austrian Space Forum led an international Mars analog mission in the Dhofar region, Oman. Directed by a Mission Support Center in Austria, a small field crew conducted experiments preparing for future human Mars missions in the fields of engineering, planetary surface operations, astrobiology, geophysics/geology and life sciences.

In summer 2017 our ScanMars experiment has been selected to be part of the mission. ScanMars uses a Ground Penetrating Radar to investigate the subsurface by transmitting and receiving radio-wave impulses into the ground. The expected results of the experiment are geophysical images of the underground structures, material differences and the presence of groundwater[1].

Here we present the ScanMars experiment and its results from AMADEE-18 analog mission to Mars.

1 Introduction

The ScanMars experiment onboard AMADEE-18 consists in a Zond Ground Penetrating Radar. We choose the 500 MHz operative frequency as a good compromise between resolution, penetration depth and maneuverability. The experiment can be divided in three phases: the training, the scientific campaign and the scientific synthesis.

2. Training Phase

The most different element from ScanMars a common radar fieldwork is undoubtedly the fact that the data was going to be acquired by the analog astronauts and not the



Figure 1: Analog astronauts during the training at OEWF headquarters in Innsbruck, Austria.

scientists. Thus the astronauts' training becomes a critical part of the experiment. The challenge of this phase resides in the fact that the analog astronauts have a background which is not specifically trimmed on the experiment, and that they have to acquire a large quantity of diverse information during the training (Figure 1).

3. Scientific campaign

The training efforts have turned into the ScanMars scientific campaign in the field in the Dhofar (Figure 2). The experiment explored four different sites with slight different geologic characteristics. The analog astronauts acquired about 1400 meters of profiles. A total of about 70000 radar echoes have been recorded and returned to the scientific data archive.

4. Results

Figure 3 shows one of the about 25 radargrams which have been acquired in the field. The data quality is very



Figure 2: The analog astronauts during ScanMars experiment data acquisition. The radar sledge is being pulled along the planned profile.




Figure 3: The radargram is the visual representation of the radar echoes recorded in the field, representing distance along the profile (horizontal) versus time delay (vertical).

good and the radar echoes show extremely well geologic features typical of the wadi riverbed. The penetration depth of the signals is about 5 meters, and in some case we can extract deeper informations.

5. Conclusions

The ScanMars experiment can be considered successful not just because of the data returned but more importantly for the volume of new experience created among the scientific team, the operation rooms and the field crew. A good teamwork has been necessary to face the problems which are difficult, when not impossible, to foresee beforehand. With ScanMars and AMADEE-18 in general we are understanding the different aspects of scientific exploration of distant worlds, developing strategies and workflows which will be the building blocks of the future human planetary missions.

Acknowledgements

Thanks to the AMADEE-18 field crew, whose efforts made this scientific experiment a successful one. We are grateful to the Italian Radio Amateur Association (ARI) which assisted in laboratory measurements and system optimization of the ScanMars radar system.

References

- [1] Annan, A. and Davis, J. L., 1989, Ground penetrating radar for high-resolution mapping of soil and rock stratigraphy. *Geophysical Prospecting* 37, 531-551
- [2] Groemer, G. et al., 2016, 47th LPSC, abstract #1903.
- [3] Rossi, A. P., 2018 Augmented field Geology and Geophysics for Planetary Analogues, European Geophysical Union, Vienna.
- [4] Snook and Mendell, 2004, The need for analogue missions in scientific human and robotic planetary exploration. 35th LPSC, abstract #2130